

Hecke Algebras and the Schlichting completion for discrete quantum groups

Roland Vergnioux

joint work with A. Skalski et C. Voigt

Université de Caen

Waterloo, May 27th, 2022

Outline

- 1 Introduction
- 2 Hecke algebras for discrete quantum groups
 - Structure of the quotient space
 - Hecke algebra
- 3 The Schlichting Completion
 - Construction of the completion
 - Application to Hecke operators
 - Exemples

Introduction

Let $\Lambda \subset \Gamma$ be a quantum subgroup of a discrete quantum group. The von Neumann algebra of Hecke operators is the commutant of the quasi-regular representation: $\mathcal{L}(\Gamma, \Lambda) = B(\ell^2(\Gamma/\Lambda))^\Gamma$.

Questions:

- ❶ (combinatorial) description of $\mathcal{L}(\Gamma, \Lambda)$?
- ❷ modular properties of the canonical state $(\delta_\Lambda | \cdot \delta_\Lambda)$?
- ❸ analytical properties of this von Neumann algebra...

In the classical case ❶ is achieved using a dense subalgebra $\mathcal{H}(\Gamma, \Lambda) \simeq c_c(\Lambda \backslash \Gamma / \Lambda)$ with convolution product. Quantum case:

- ❹ construction and description of $\Gamma/\Lambda, \Lambda \backslash \Gamma / \Lambda$?
- ❺ boundedness of the action of $c_c(\Lambda \backslash \Gamma / \Lambda)$ on $\ell^2(\Gamma/\Lambda)$?

For ❸, ❺ we construct the Schlichting completion (\mathbb{G}, \mathbb{H}) of (Γ, Λ) .

→ construction of new locally compact quantum groups.

Outline

- 1 Introduction
- 2 Hecke algebras for discrete quantum groups
 - Structure of the quotient space
 - Hecke algebra
- 3 The Schlichting Completion
 - Construction of the completion
 - Application to Hecke operators
 - Exemples

Quantum groups

An **algebraic quantum group** [Van Daele] \mathbb{G} is given by

- a (non-unital) $*$ -algebra $\mathcal{O}_c(\mathbb{G})$,
- a coproduct $\Delta : \mathcal{O}_c(\mathbb{G}) \rightarrow \mathcal{M}(\mathcal{O}_c(\mathbb{G}) \odot \mathcal{O}_c(\mathbb{G}))$

with some axioms, in particular:

- $\Delta(\mathcal{O}_c(\mathbb{G}))(\mathcal{O}_c(\mathbb{G}) \otimes 1) \subset \mathcal{O}_c(\mathbb{G}) \odot \mathcal{O}_c(\mathbb{G})$,
- left integral: $\varphi : \mathcal{O}_c(\mathbb{G}) \rightarrow \mathbb{C}$ s.t. $(\text{id} \otimes \varphi)((a \otimes 1)\Delta(b)) = \varphi(b)a$.

Commutative case: there exists a locally compact group G with a compact-open $H \subset G$ s.t.

$$\mathcal{O}_c(\mathbb{G}) = \{f \in C_c(G) \mid \dim \text{Vect}(H \cdot f) < \infty\}.$$

Quantum groups

An **algebraic quantum group** [Van Daele] \mathbb{G} is given by

- a (non-unital) $*$ -algebra $\mathcal{O}_c(\mathbb{G})$,
- a coproduct $\Delta : \mathcal{O}_c(\mathbb{G}) \rightarrow \mathcal{M}(\mathcal{O}_c(\mathbb{G}) \odot \mathcal{O}_c(\mathbb{G}))$

with some axioms, in particular:

- $\Delta(\mathcal{O}_c(\mathbb{G}))(\mathcal{O}_c(\mathbb{G}) \otimes 1) \subset \mathcal{O}_c(\mathbb{G}) \odot \mathcal{O}_c(\mathbb{G})$,
- left integral: $\varphi : \mathcal{O}_c(\mathbb{G}) \rightarrow \mathbb{C}$ s.t. $(\text{id} \otimes \varphi)((a \otimes 1)\Delta(b)) = \varphi(b)a$.

Discrete case: $\mathcal{O}_c(\Gamma) \simeq \bigoplus_{\alpha \in I(\Gamma)} B(H_\alpha)$, $\dim H_\alpha < \infty$.

The existence of integrals $\varphi := h_L, h_R$ is automatic.

We put $c_c(\Gamma) = \mathcal{O}_c(\Gamma)$, $\ell^\infty(\Gamma) = M(c_c(\Gamma))$.

Denote $p_\alpha = \text{id}_{H_\alpha}$, $a_\alpha = p_\alpha a \in B(H_\alpha)$ for $a \in \ell^\infty(\Gamma)$, $\alpha \in I(\Gamma)$.

Δ endows the C^* -category $\text{Corep}(\Gamma) := \text{Rep}(c_c(\Gamma))$ with a tensor structure:

$v \otimes w := (v \otimes w) \circ \Delta$. Underlying combinatorial data: spaces $\text{Hom}_\Gamma(\alpha, \beta \otimes \gamma)$ for $\alpha, \beta, \gamma \in I(\Gamma)$.

Subgroups of discrete quantum groups

A (quantum) **subgroup** $\Lambda \subset \Gamma$ is given by $\ell^\infty(\Lambda) \simeq p_\Lambda \ell^\infty(\Gamma)$ for some central proj. $p_\Lambda \in \ell^\infty(\Gamma)$ s.t. $\Delta(p_\Lambda)(1 \otimes p_\Lambda) = p_\Lambda \otimes p_\Lambda = \Delta(p_\Lambda)(p_\Lambda \otimes 1)$. We have $I(\Lambda) \subset I(\Gamma)$, $\text{Corep}(\Lambda) \subset \text{Corep}(\Gamma)$, $p_\Lambda = \sum_{\alpha \in I(\Lambda)} p_\alpha$.

Quotient spaces.

- Quantum: $\ell^\infty(\Gamma/\Lambda) = \ell^\infty(\Gamma)^\wedge = \{a \in \ell^\infty(\Gamma) \mid (1 \otimes p_\Lambda)\Delta(a) = a \otimes p_\Lambda\}$. We have $\Delta(\ell^\infty(\Gamma/\Lambda)) \subset \ell^\infty(\Gamma) \bar{\otimes} \ell^\infty(\Gamma/\Lambda)$.
- Categorical: $\text{Corep}(\Gamma/\Lambda) := \text{Rep}(\ell^\infty(\Gamma/\Lambda))$ is a left- $\text{Corep}(\Gamma)$ -module category with restriction functor $\text{Corep}(\Gamma) \rightarrow \text{Corep}(\Gamma/\Lambda)$.
- Classical: $I(\Gamma)/\Lambda = I(\Gamma)/\sim$ where $\alpha \sim \beta \Leftrightarrow \exists \lambda \in I(\Lambda) \quad \beta \subset \alpha \otimes \lambda$.

Some notation:

- $v_\Lambda \in \text{Corep}(\Lambda)$, Λ -isotypical component of $v \in \text{Corep}(\Gamma)$,
- $\kappa_\alpha = \dim_q(\bar{\alpha} \otimes \alpha)_\Lambda$ for $\alpha \in I(\Gamma)$,
- $[\alpha]$ the class of α in $I(\Gamma)/\Lambda$ (or $\Lambda \setminus I(\Gamma)$),
- $p_{[\alpha]} = \sum_{\beta \in [\alpha]} p_\beta \in \ell^\infty(\Gamma/\Lambda)$.

Description of the quotient space

Theorem

- We have $\ell^\infty(\Gamma/\Lambda) = \ell^\infty - \bigoplus_{[\alpha]} p_{[\alpha]} \ell^\infty(\Gamma/\Lambda)$.
- We have $p_{[\alpha]} \ell^\infty(\Gamma/\Lambda) \simeq B(H_\alpha)'_\Lambda \cap B(H_\alpha)$ canonically.
- More generally $\text{Hom}_{\Gamma/\Lambda}(\alpha, \beta) = B(H_\alpha, H_\beta)_\Lambda$.
- For $a \in \ell^\infty(\Gamma/\Lambda)$ we have $a = \sum_{[\alpha]} \kappa_\alpha^{-1}(h_R \otimes \text{id})[(S^{-1}(a_\alpha) \otimes 1)\Delta(p_\Lambda)]$.

Denote $c_c(\Gamma/\Lambda) = \bigoplus p_{[\alpha]} \ell^\infty(\Gamma/\Lambda)$, $c(\Gamma/\Lambda) = \prod p_{[\alpha]} \ell^\infty(\Gamma/\Lambda)$.

Analogue of the counting measure on Γ/Λ :

Corollary

$c_c(\Gamma/\Lambda)$ admits a (unique) positive, Γ -invariant faithful form given by

$$\mu(a) = \sum_{[\alpha]} \kappa_\alpha^{-1} h_L(a_\alpha).$$

The Hecke algebra

Definition

For $a \in c_c(\Gamma/\Lambda)$, $b \in c_c(\Lambda \setminus \Gamma)$ define

$$a * b = (\text{id} \otimes \mu)[\Delta(a)(1 \otimes S(b))] = (\mu S \otimes \text{id})[\Delta(b)(S^{-1}(a) \otimes 1)] \in c(\Gamma)$$

and $a^\# = S(a^*) \in c_c(\Lambda \setminus \Gamma)$.

Classical case: $\forall \alpha \dim_q(\alpha) = 1$, $\kappa_\alpha = 1$. We recover the formula

$$(a * b)(g) = \sum_{[h] \in \Gamma/\Lambda} a(gh)b(h^{-1}).$$

Proposition-Definition

$\mathcal{H}(\Gamma, \Lambda) := c_c(\Gamma/\Lambda) \cap c_c(\Lambda \setminus \Gamma)$ is an involutive algebra for $*$ and $\#$, with unit p_Λ , stable under σ_t^R , σ_t^L and τ_t .

The Hecke algebra

Proposition-Definition

$\mathcal{H}(\Gamma, \Lambda) := c_c(\Gamma/\Lambda) \cap c_c(\Lambda \setminus \Gamma)$ is an involutive algebra for $*$ and \sharp , with unit p_Λ , stable under σ_t^R , σ_t^L and τ_t .

NB. $c_c(\Gamma/\Lambda) \cap c_c(\Lambda \setminus \Gamma)$ is also a (possibly degenerate) sub- $*$ -algebra of $\ell^\infty(\Gamma)$. Denote, for $\tau \in \Lambda \setminus I(\Gamma)/\Lambda$:

$$L(\tau) = \#\{\alpha \in \Lambda \setminus I(\Gamma) \mid \alpha \subset \tau\}, \quad R(\tau) = \#\{\alpha \in I(\Gamma)/\Lambda \mid \alpha \subset \tau\}.$$

Proposition-Definition

We say that (Γ, Λ) is a Hecke pair if $c_c(\Gamma/\Lambda) \cap c_c(\Lambda \setminus \Gamma) \subset \ell^\infty(\Gamma)$ is non degenerate $\Leftrightarrow \forall \tau \in \Lambda \setminus I(\Gamma)/\Lambda \quad L(\tau) < \infty$.

Examples: Λ finite or finite index. Normal case: if $c_c(\Gamma/\Lambda) = c_c(\Lambda \setminus \Gamma)$, $\mathcal{H}(\Gamma, \Lambda)$ is the convolution algebra of the quotient quantum group Γ/Λ .

Hecke Operators

Recall that $c_c(\Gamma/\Lambda)$ is endowed with a left Γ -action.

Proposition

Let (Γ, Λ) be a Hecke pair. We have an isomorphism

$$\mathcal{H}(\Gamma, \Lambda) \rightarrow \text{End}(c_c(\Gamma/\Lambda))^\Gamma, a \mapsto T(a) := (\cdot * a)$$

with inverse $(T \rightarrow T(p_\Lambda))$. Moreover $(x | T(a)y) = (T(a^\#)x | y)$ for the scalar product associated with μ .

Theorem

Let (Γ, Λ) be a Hecke pair. $T(a)$ is bounded on $\ell^2(\Gamma/\Lambda)$ for all $a \in \mathcal{H}(\Gamma, \Lambda)$ **iff** we have $(RT) : \kappa_\gamma \leq C_\beta \kappa_\alpha$ for all $\gamma \subset \alpha \otimes \beta$.

Hecke Operators

Theorem

Let (Γ, Λ) be a Hecke pair. $T(a)$ is bounded on $\ell^2(\Gamma/\Lambda)$ for all $a \in \mathcal{H}(\Gamma, \Lambda)$ **iff** we have (RT) : $\kappa_\gamma \leq C_\beta \kappa_\alpha$ for all $\gamma \subset \alpha \otimes \beta$.

This property is not satisfied by all inclusions $\Lambda \subset \Gamma$.

By restriction to $\ell^\infty(\Gamma/\Lambda)$ we have also $\alpha \in \text{Corep}(\Gamma/\Lambda)$. Denote $\tilde{\alpha}$ is the image of α in the Grothendieck ring $\mathbb{Z}[\Gamma/\Lambda]$. If Γ is unimodular:

$$\kappa_\alpha = \dim_q(\tilde{\alpha} \otimes \alpha)_\Lambda = \|\tilde{\alpha}\|_2^2.$$

Exercise

Prove in $\text{Corep}(\Gamma/\Lambda)$ that (RT) is satisfied if (Γ, Λ) is a Hecke pair.

In the sequel we will see an analytical proof.

Modular properties

Canonical state on $\mathcal{H}(\Gamma, \mathbb{A})$: $\omega = \epsilon = (p_{\mathbb{A}} | T(\cdot)p_{\mathbb{A}})$. It is faithful.

Proposition-Definition

Let $\nabla \in c(\mathbb{A} \setminus \Gamma / \mathbb{A})$ unique such that $\mu S(a) = \mu(\nabla a)$ for all $a \in \mathcal{H}(\Gamma, \mathbb{A})$. Then $\theta_t : a \mapsto \sigma_t^R(\nabla^{it} a)$ is a group of \sharp -automorphisms of $\mathcal{H}(\Gamma, \mathbb{A})$ and ω is θ -KMS.

Theorem

Assume \mathbb{A} is unimodular. Then $\nabla_{\alpha} = (\tilde{L}(\llbracket \alpha \rrbracket) / \tilde{R}(\llbracket \alpha \rrbracket)) F_{\alpha}^2$ where the F_{α} are Woronowicz' modular matrices, and for $\tau = \llbracket \alpha \rrbracket \in \mathbb{A} \setminus I(\Gamma) / \mathbb{A}$:

$$\begin{aligned}\tilde{L}(\tau) &= \sum_{[\delta] \in \mathbb{A} \setminus I(\Gamma), [\delta] \subset \tau} (\dim_q \delta)^2 / \kappa_{\bar{\delta}} \\ \tilde{R}(\tau) &= \sum_{[\delta] \in I(\Gamma) / \mathbb{A}, [\delta] \subset \tau} (\dim_q \delta)^2 / \kappa_{\delta}.\end{aligned}$$

There is also a more involved formula when \mathbb{A} is not unimodular...

Outline

- 1 Introduction
- 2 Hecke algebras for discrete quantum groups
 - Structure of the quotient space
 - Hecke algebra
- 3 **The Schlichting Completion**
 - Construction of the completion
 - Application to Hecke operators
 - Exemples

The C^* -Hopf algebra

Classical reminder. Consider $\pi : \Gamma \rightarrow \text{Bij}(\Gamma/\Lambda)$ by left translations.

Define $G = \overline{\pi(\Gamma)}$. If (Γ, Λ) is a Hecke pair, it is a locally compact group and $H = \overline{\pi(\Lambda)}$ is compact open.

But $\text{Bij}(\Gamma/\Lambda)$ has no good quantum analogue...

The C^* -Hopf algebra

Classical reminder. Consider $\pi : \Gamma \rightarrow \text{Bij}(\Gamma/\Lambda)$ by left translations.

Define $G = \overline{\pi(\Gamma)}$. If (Γ, Λ) is a Hecke pair, it is a locally compact group and $H = \overline{\pi(\Lambda)}$ is compact open.

But $\text{Bij}(\Gamma/\Lambda)$ has no good quantum analogue...

Definition

We put $\mathcal{O}_c(G) = \text{alg}\langle a * b, a \in c_c(\Gamma/\Lambda), b \in c_c(\Lambda \backslash \Gamma) \rangle \subset c(\Gamma)$.

$$C_0(G) = C^*\langle a * b, a \in c_c(\Gamma/\Lambda), b \in c_c(\Lambda \backslash \Gamma) \rangle \subset \ell^\infty(\Gamma).$$

Classical case: $\Gamma \rightarrow G$ induces $C_0(G) \subset \ell^\infty(\Gamma)$.

If $a = \mathbb{1}_{[r]}$, $b = \mathbb{1}_{[s]}$ then $a * b = \mathbb{1}_{\{g \mid g[s]^{-1} = [r]\}}$.

The C^* -Hopf algebra

Definition

We put $\mathcal{O}_c(\mathbb{G}) = \text{alg}\langle a * b, a \in c_c(\Gamma/\Lambda), b \in c_c(\Lambda \setminus \Gamma) \rangle \subset c(\Gamma)$.
 $C_0(\mathbb{G}) = C^*\langle a * b, a \in c_c(\Gamma/\Lambda), b \in c_c(\Lambda \setminus \Gamma) \rangle \subset \ell^\infty(\Gamma)$.

Classical case: $\Gamma \rightarrow G$ induces $C_0(\mathbb{G}) \subset \ell^\infty(\Gamma)$.

If $a = \mathbb{1}_{[r]}$, $b = \mathbb{1}_{[s]}$ then $a * b = \mathbb{1}_{\{g \mid g[s]^{-1}=[r]\}}$.

Theorem

If (Γ, Λ) is a Hecke pair we have

$$\Delta(\mathcal{O}_c(\mathbb{G}))(1 \otimes \mathcal{O}_c(\mathbb{G})) = \mathcal{O}_c(\mathbb{G}) \odot \mathcal{O}_c(\mathbb{G}) = \Delta(\mathcal{O}_c(\mathbb{G}))(\mathcal{O}_c(\mathbb{G}) \otimes 1).$$

$\mathcal{O}_c(\mathbb{G})$ is a multiplier Hopf algebra.

$C_0(\mathbb{G})$ is a bisimplifiable Hopf C^* -algebra.

The Haar weights

Corollary-Definition

$C(\mathbb{H}) := p_{\wedge} C_0(\mathbb{G})$ is a Hopf C^* -algebra (with unit p_{\wedge}).

Hence it admits a Haar state h .

We have $c_c(\Gamma/\wedge) = c_c(\mathbb{G}/\mathbb{H})$ in $\ell^\infty(\Gamma)$.

Corollary-Definition

$\varphi := \mu(\text{id} \otimes hp_{\wedge})\Delta$ is an integral on $\mathcal{O}_c(\mathbb{G})$. \mathbb{G} is an algebraic quantum group, hence a locally compact quantum group.

Proposition

If the action of Γ on Γ/\wedge is faithful and \wedge is infinite, \mathbb{G} is non-discrete.

The Hecke Algebra

Let (\mathbb{G}, \mathbb{H}) be the Schlichting completion of a Hecke pair (Γ, Λ) .

Recall that \mathbb{H} is compact and put

$$c_c(\mathbb{G}/\mathbb{H}) := \mathcal{O}_c(\mathbb{G})^{\mathbb{H}} \subset \mathcal{O}_c(\mathbb{G}) \text{ and } \ell^2(\mathbb{G}/\mathbb{H}) = \overline{c_c(\mathbb{G}/\mathbb{H})} \subset L^2(\mathbb{G}).$$

Proposition

We have ${}^{\mathbb{H}}\mathcal{O}_c(\mathbb{G})^{\mathbb{H}} \simeq \text{End}(c_c(\mathbb{G}/\mathbb{H}))^{\mathbb{G}}$, $b \mapsto T'(b) := (\cdot * b)$, using the convolution product of $\mathcal{O}_c(\mathbb{G})$.

By construction of (\mathbb{G}, \mathbb{H}) we have

$$\text{End}(c_c(\mathbb{G}/\mathbb{H}))^{\mathbb{G}} = \text{End}(c_c(\Gamma/\Lambda))^{\Gamma} \text{ and } \ell^2(\mathbb{G}/\mathbb{H}) \simeq \ell^2(\Gamma/\Lambda).$$

Since the operators $T'(b)$ arise from the right regular repr. of \mathbb{G} we get:

Corollary

The Hecke operators $T(a)$, for $a \in \mathcal{H}(\Gamma, \Lambda)$, are bounded on $\ell^2(\Gamma/\Lambda)$.

HNN Extensions

Fixe $\Lambda_{\pm 1} \subset \Gamma_0$ with an isomorphism $\theta : \Lambda_1 \rightarrow \Lambda_{-1}$.

Consider $\Gamma = \text{HNN}(\Gamma_0, \theta)$ [Fima 2013]. $\text{Corep}(\Gamma)$ is generated by $\text{Corep}(\Gamma_0)$ and a 1-dimensional w such that $w^\epsilon \otimes v \otimes w^{-\epsilon} = \theta_*^\epsilon(v)$ for $v \in \text{Corep}(\Lambda_\epsilon)$.

Proposition

- If $\Lambda_{\pm 1}$ have finite index in Γ_0 and are different from Γ_0 , then $\Gamma_0 \subset \Gamma$ is almost normal, not normal, of infinite index.
- If $\bigcap_{k \in \mathbb{Z}} \text{Dom } \theta^k = \{1\}$ the action of Γ on Γ/Γ_0 is faithful.
- We have $\nabla_w = (\tilde{L}(\llbracket w \rrbracket) / \tilde{R}(\llbracket w \rrbracket)) p_w$ with

$$\tilde{L}(\llbracket w \rrbracket) = \sum_{[\delta] \in \Lambda_1 \setminus I(\Gamma_0)} \dim_q(\delta \otimes \bar{\delta}) / \dim_q(\delta \otimes \bar{\delta})_{\Lambda_1},$$

$$\tilde{R}(\llbracket w \rrbracket) = \sum_{[\delta] \in I(\Gamma_0) / \Lambda_{-1}} \dim_q(\bar{\delta} \otimes \delta) / \dim_q(\bar{\delta} \otimes \delta)_{\Lambda_{-1}}.$$

HNN Extensions

Proposition

- If $\Lambda_{\pm 1}$ have finite index in Γ_0 and are different from Γ_0 , then $\Gamma_0 \subset \Gamma$ is almost normal, not normal, of infinite index.
- If $\bigcap_{k \in \mathbb{Z}} \text{Dom } \theta^k = \{1\}$ the action of Γ on Γ/Γ_0 is faithful.
- We have $\nabla_w = (\tilde{L}(\llbracket w \rrbracket) / \tilde{R}(\llbracket w \rrbracket)) \rho_w$ with

$$\tilde{L}(\llbracket w \rrbracket) = \sum_{[\delta] \in \Lambda_1 \setminus I(\Gamma_0)} \dim_q(\delta \otimes \bar{\delta}) / \dim_q(\delta \otimes \bar{\delta})_{\Lambda_1},$$

$$\tilde{R}(\llbracket w \rrbracket) = \sum_{[\delta] \in I(\Gamma_0) / \Lambda_{-1}} \dim_q(\bar{\delta} \otimes \delta) / \dim_q(\bar{\delta} \otimes \delta)_{\Lambda_{-1}}.$$

Example. $\Sigma_{\pm 1}$ non classical finite quantum groups.

Take $\Gamma_0 = \prod'_{k \in \mathbb{Z}^*} \Sigma_{\text{sgn}(k)}$ and $\Lambda_{\epsilon} = \prod'_{k \in \mathbb{Z}^*, k \neq \epsilon} \Sigma_{\text{sgn}(k)} \subset \Gamma_0$.

Then the Proposition applies, $\tilde{L}(\llbracket w \rrbracket) = \#\Sigma_1$, $\tilde{R}(\llbracket w \rrbracket) = \#\Sigma_{-1}$.

\mathbb{G} is a non-discrete, non-classical, non-co-classical locally compact quantum group, with non-trivial modular group if $\#\Sigma_1 \neq \#\Sigma_{-1}$.